arXiv:0706.1165v1 [physics.flu-dyn] 8 Jun 2007

On the existence of two-dimensional nonlinear steady states in plane Couette flow

F. Rincon*
Department of Applied Mathematics and Theoretical Physics, University of Cambridge,
Centre for Mathematical Sciences, Wilberforce Road, Cambridge CB3 0WA, United Kingdom

(Dated: February 1, 2008)

The problem of two-dimensional steady nonlinear dynamics in plane Couette flow is revisited using homotopy from either plane Poiseuille flow or from plane Couette flow perturbed by a small symmetry-preserving identity operator. Our results show that it is not possible to obtain the nonlinear plane Couette flow solutions reported by Cherhabili and Ehrenstein [Eur. J. Mech. B/Fluids, 14, 667 (1995)] using their Poiseuille-Couette homotopy. We also demonstrate that the steady solutions obtained by Mehta and Healey [Phys. Fluids, 17, 4108 (2005)] for small symmetry-preserving perturbations are influenced by an artefact of the modified system of equations used in their paper. However, using a modified version of their model does not help to find plane Couette flow solution in the limit of vanishing symmetry-preserving perturbations either. The issue of the existence of two-dimensional nonlinear steady states in plane Couette flow remains unsettled.

PACS numbers: 47.20.Ft, 47.15.Fe, 47.20.Ky, 47.27.Cn, 47.27.De, 47.27.N-, 47.32.Ef

I. INTRODUCTION

Even though plane Couette flow is known to be linearly stable for all values of the Reynolds number Re, transition to turbulence is observed experimentally and numerically at quite modest values of Re $\simeq 350$ in this flow[1]. As a consequence, the quest for nonlinear instabilities has been the subject of many studies. Since there is no direct connection between the basic state and potential nonlinear branches of solutions, homotopy (continuation) from other flows subject to linear instabilities has been a very popular method to discover such branches. Exact three-dimensional solutions have been reported by Nagata[2] using homotopy from linearly unstable rotating plane Couette flow, by Clever and Busse[3] using homotopy from a convectively unstable plane Couette flow and by Waleffe[4,5] using a forcing homotopy. These three-dimensional states match very well experimental data on transition to turbulence in plane Couette flow, but other solutions, including two-dimensional ones, may well be participating in phase-space dynamics and in the transition process.

Another kind of homotopy was suggested by Milinazzo and Saffman[6] to discover two-dimensional nonlinear solutions in plane Couette flow by connecting the nonbifurcating no-slip plane Couette flow to the no-slip plane Poiseuille flow, which becomes linearly unstable at Re = 5772 (Zahn et al.[7], Herbert[8]). Starting from the travelling-wave solutions in this flow and progressively changing the original background flow from a Poiseuille to a Couette profile, Milinazzo and Saffman[6] attempted to continue these solutions to a pure Couette flow, but they failed to achieve this goal. Their lack of success may have to do with resolution: Cherhabili and Ehrenstein[9] (hereinafter CE95) used the same approach with more Fourier modes in the streamwise direction and claimed to have identified steady two-dimensional localized nonlinear solutions for the Couette flow. Cherhabili and Ehrenstein[10] subsequently computed the stability of these two-dimensional steady states and reported the discovery of three-dimensional nonlinear structures which are different from those found by Nagata[2].

Recently, Mehta and Healey[11] (hereinafter MH05) suggested yet another interesting homotopy to compute two-dimensional nonlinear states in plane Couette flow. They argued that the homotopy from Poiseuille to Couette flow was not symmetry-preserving, so that possible nonlinear states relying on the $O(2)$ symmetry of the Couette flow might not be found by continuation from flows that do not have this symmetry. Instead, they proposed a symmetry-preserving homotopy based on small perturbations of the linearization of the Couette operator that causes the least stable eigenvalues of the plane Couette flow to become unstable. Doing so, they hoped to be able to track the nonlinear branch associated with this bifurcation point down to zero perturbation and to obtain two-dimensional nonlinear states for the unperturbed plane Couette flow. For small perturbation parameters, they discovered steady solutions looking similar to the solutions of CE95. MH05 notably describe the numerical appearance of so-called "frozen-waves" with phase speed $c = 0$ in perturbed Couette flow starting from bifurcating travelling-wave solutions at Re > 390. CE95 had also noticed the numerical appearance of solutions with $c = 0$ in their calculations. However, these solutions cannot be followed down to zero perturbation (exact plane Couette flow), which raises the questions of the existence of the solutions reported by CE95 and of the connections between their findings and those of MH05.

Considering that the case is quite unclear, we attempt to address some of these issues in this study. We first revisit the Poiseuille-Couette homotopy problem and ar-

---

*Electronic address: F.Rincon@damtp.cam.ac.uk

gue that the solutions obtained by CE95 are spurious. Then, we clarify the nature of the steady solutions found by MH05 at large Re for small values of their perturbation parameter ε. We show that a direct connection exists between travelling and steady solutions for the perturbed flow and demonstrate that their high Reynolds number steady solutions are influenced by an artifact in the model that introduces a mean flow eigenmode. We obtain solutions for small perturbations using a revised symmetry-preserving homotopy and show that they cannot be continued to non-perturbed plane Couette flow either. Our results therefore suggest that it is not possible to identify two-dimensional nonlinear solutions in plane Couette flow using these two particular methods.

The paper is organized as folllows. In Sec. II, we formulate the physical problem, describe the underlying symmetries and present the numerical techniques used to compute nonlinear branches of solutions. Sections III and IV describe our efforts to obtain steady two-dimensional nonlinear plane Couette solutions using either the Poiseuille-Couette homotopy with the same procedure as CE95 or the symmetry-preserving homotopy of MH05. A short discussion of the results (Sec. V) concludes the paper.

II. EQUATIONS, SYMMETRIES, NUMERICS

A. Physical model and background state

In the following, an incompressible fluid with kinematic viscosity ν and constant density ρ taken to be unity is considered. The plane shear flows investigated here are independent of the streamwise coordinate x and are bounded by two rigid walls in the shearwise y direction, with $-d < y < d$, where d is chosen as length unit. The x direction is taken to be periodic with spatial period $L_x = 2\pi d/\alpha$. A reference velocity U_o is defined and a mean streamwise pressure gradient $-2\nu U_o(1-\eta)/d^2$ and mean streamwise wall velocities $U_B(y=\pm 1) = \pm\eta U_o$ are imposed in order to maintain the following generic (nondimensional) streamwise oriented background shear flow:

$$U_B(y) = (1-\eta)\left(1-y^2\right) + \eta\, y\ . \tag{1}$$

Here, η is a nondimensional parameter that allows continuation from plane Poiseuille flow ($\eta = 0$) to plane Couette flow ($\eta = 1$). The Reynolds number of the flow, based on U_o and half of the channel width, reads $\mathrm{Re} = (U_o d)/\nu$.

B. Equations

The full non-dimensional velocity field, denoted by $\boldsymbol{U}$, is decomposed into the original streamwise background flow $(U_B, 0)$ and two-dimensional velocity perturbations denoted by $\boldsymbol{u} = (u, v)$. The corresponding total pressure is $P = P_B + p$, where p is a pressure perturbation, and the nondimensional equations governing the evolution of $\boldsymbol{U}$ and P are the incompressible Navier-Stokes equations

$$\partial_t \boldsymbol{U} + \boldsymbol{U}\cdot\boldsymbol{\nabla}\boldsymbol{U} = -\boldsymbol{\nabla}P + \frac{1}{\mathrm{Re}}\Delta\boldsymbol{U}\ , \quad \boldsymbol{\nabla}\cdot\boldsymbol{U} = 0. \tag{2}$$

They are supplemented by periodic boundary conditions in x and by no-slip boundary conditions in y for u and v:

$$u(x,-1,t) = u(x,1,t) = v(x,-1,t) = v(x,1,t) = 0\ , \tag{3}$$

For numerical purposes, a Fourier decomposition is used in the x direction and a scalar equation for the shearwise velocity v (or equivalently for a stream function ψ) governing the evolution of the $k_x \neq 0$ modes is solved:

$$\left(\partial_t - \frac{1}{\mathrm{Re}}\Delta\right)\Delta v + \boldsymbol{P}_v\cdot(\boldsymbol{U}\cdot\boldsymbol{\nabla}\boldsymbol{U}) = 0\ , \tag{4}$$

where $\boldsymbol{P}_v = -\boldsymbol{e}_y\cdot\boldsymbol{\nabla}\times\boldsymbol{\nabla}\times(\cdot)$. An auxiliary equation describing the evolution of the mean streamwise velocity field must be solved simultaneously for completeness:

$$\mathrm{Re}\left(\partial_t\overline{u} + \partial_y\overline{uv}\right) = \partial_y^2\overline{u}\ , \tag{5}$$

where overbars denote x-averaged quantities.

C. Symmetries

Depending on the value of η, this set of equations has different symmetries. For $\eta \neq 1$, the only symmetry is the translation symmetry $SO(2)$

$$SO(2) : (x,y) \to (x+a,y)\ , \quad (u,v,p) \to (u,v,p) \tag{6}$$

with $a \in [0, 2\pi/\alpha]$. For plane Couette flow ($\eta = 1$), an extra (point) reflection symmetry Z^2 is present,

$$Z^2 : (x,y) \to (-x,-y)\ , \quad (u,v,p) \to (-u,-v,p)\ . \tag{7}$$

In that case, the full flow has the more general $O(2)$ symmetry.

D. A symmetry preserving homotopy

An extra symmetry-preserving artificial term proportional to $\varepsilon\boldsymbol{u}$, where ε is a small non-dimensional parameter, can be added to the r.h.s. of the original Navier-Stokes equations (2) in order to cause the least stable eigenvalue to become unstable and to perform the symmetry-preserving homotopy mentioned in the introduction. This technique is described in detail in MH05.

E. Numerics

The exact two-dimensional nonlinear steady and travelling solutions of Eqs. (3)-(5) presented in the next sections have been obtained thanks to a nonlinear continuation code[12] similar to that of Waleffe[5] and Wedin and

Kerswell[13]. Chebyshev collocation on a Gauss-Lobatto grid in the shearwise direction and a Fourier representation in the x direction are used. Clamped boundary conditions are implemented using differentiation matrices computed with the DMSuite package[14]. Nonlinear terms are computed in physical space and steady nonlinear equations are solved in spectral-physical-spectral space using Newton iteration. A direct LAPACK solver is called at each Newton iteration and dealiasing is performed in the x-direction. Pseudo-arclength continuation[15] is implemented to follow branches of solutions. The code has been tested quantitatively by solving the classical problem of subcritical continuation of two-dimensional travelling waves in no-slip plane Poiseuille flow[7,8]. The full three-dimensional version (not used here) also quantitatively reproduces the continuation diagrams of three-dimensional nonlinear solutions obtained by Waleffe[5] in no-slip, non-rotating plane Couette flow[12].

III. POISEUILLE-COUETTE HOMOTOPY

The Poiseuille-Couette homotopy was first considered in order to compute the two-dimensional nonlinear solutions reported by CE95 in plane Couette flow. No particular assumption regarding the symmetries of the solutions was made in order to perform these computations and the numerical method employed here copes with solutions with either zero or non-zero phase speed (the transformation $\partial_t = -c\,\partial_x$, where c is the phase speed, is used in the latter case). All our attempts proved unsuccessful. We simply followed their procedure and performed continuation with respect to η, starting from two-dimensional nonlinear travelling wave solutions in Poiseuille flow ($\eta = 0$) at $\mathrm{Re} = 15993$, which corresponds to the first case considered by CE95. Continuation curves for mixed Poiseuille-Couette flow are shown in Fig. 1. For the maximum affordable resolution and $\alpha = 1$, a turning point is found at $\eta \simeq 0.23$, therefore it proves impossible to isolate solutions for plane Couette flow. Since CE95 found Couette solutions for small α only (albeit for smaller Re), we tried to perform further continuation with respect to α for various values of η and different resolutions including the maximum resolution used by CE95 (corresponding to $N_x = 52$ in our code). The results, presented in Fig. 2, show that it is not possible to identify plane Couette flow solutions for small values of α either. The nose of the surface of solutions in the $\alpha - \eta -$ wall shear rate space is located at $\eta \simeq 0.5$ and $\alpha \simeq 0.2$ for $\mathrm{Re} = 15993$, even for rather large streamwise resolution. This result is similar to Fig. 1 of CE95, who presented low streamwise resolution solutions down to $\eta = 0.45$ for $\mathrm{Re} = 15993$. As already acknowledged in CE95, it is clear from the figure that there are convergence issues at low α for such a high value of Re, even with a resolution such as $(N_x, N_y) = (52, 48)$. We notably encountered difficulties performing the low-α continuation at $\eta = 0.45$. The same procedure was consequently attempted at the less resolution-demanding $\mathrm{Re} = 7003$ at which CE95 found their Couette flow solutions. The corresponding continuation curves for mixed Poiseuille-Couette flow are also shown in Fig. 1. This time, a turning point is found at $\eta \simeq 0.14$ for $\alpha = 1$. Further continuations with respect to α for various values of η and the same maximum resolution as the one used by CE95 are represented in Fig. 3 and show that it is not possible to identify plane Couette flow solutions for small values of α either. The nose of the surface of solutions in the $\alpha - \eta -$wall shear rate space is clearly located at $\eta \simeq 0.14$ and $\alpha \simeq 1$ for $\mathrm{Re} = 7003$.The same procedure was finally attempted for even smaller values of Re and led to similar negative results.



FIG. 1: Continuation of two-dimensional solutions with $\alpha = 1$ in mixed Poiseuille-Couette flow (1) as a function of η, for $\mathrm{Re} = 15993$ (black curves with turning points at $\eta \simeq 0.23$ and $\mathrm{Re} = 7003$ (blue online, with turning points at $\eta \simeq 0.13$). Full line: $(N_x, N_y) = (52, 48)$, dashed line (almost undistinguishable from full line): $(N_x, N_y) = (32, 48)$, dotted line: $(N_x, N_y) = (6, 48)$. The non-dimensional wall shear rate $|\partial\overline{U}/\partial y|(y = \pm 1)$ has been used as a measure of nonlinearity (for the laminar flow, $|\partial\overline{U}/\partial y|(\pm 1) = 1$). The "gap" at $\eta \simeq 0$ results from the existence of a turning point close to $\eta \simeq 0.02$.

The main trend is therefore that it seems possible to reach solutions with lower η with increasing Re, but it is clear from our results that no $\eta = 1$ solution can be found for the range of Re investigated by CE95. Since only very little information is given on their precise procedure to obtain two-dimensional solutions, it is actually very hard to figure out where the problem could come from in their results. We originally thought that aliasing errors may be present in their calculations, but since the numerical method used by CE95 is fully spectral, their results should not in principle suffer from this problem, at least in the streamwise direction. One possibility would be that their error tolerance for the Newton algorithm was not small enough. In the present calculations, convergence was considered to be achieved for an error smaller than 10^{-7} in energy and a precision of 10^{-9} was reached



FIG. 2: Continuation of two-dimensional solutions (Re $=$ 15993) in mixed Poiseuille-Couette flow (1) as a function of α for $\eta = 0.17$ (black curves with the largest α values), $\eta = 0.23$ (blue online, curves centered on $\alpha \simeq 0.5$), $\eta = 0.45$ (red online, narrow curves at low α. Full line: $(N_x, N_y) = (52, 48)$, dashed line: $(N_x, N_y) = (32, 48)$, dotted line:$(N_x, N_y) = (6, 48)$. Solutions for $N_x = 6$ do not extend to $\eta = 0.45$ and the corresponding red dotted curve at $\alpha \simeq 0.2$ is for $\eta = 0.4$. Note the convergence problems at low α for $(N_x, N_y) = (32, 48)$ and $(N_x, N_y) = (52, 48)$.



FIG. 3: Continuation of two-dimensional solutions (Re $=$ 7003) in mixed Poiseuille-Couette flow (1) as a function of α for $\eta = 0.05$ (black, family of large curves), $\eta = 0.12$ (blue online, family of medium size curves), $\eta = 0.132$ (red online, family of small curves corresponding to the "nose" of the surface of solutions). Full line: $(N_x, N_y) = (52, 48)$, dashed line (undistinguishable from full line curves): $(N_x, N_y) = (32, 48)$, dotted line:$(N_x, N_y) = (6, 48)$. The results for $(N_x, N_y) = (52, 48)$ presented here are also undistinguishable from those (not shown here) obtained for the same resolution $(N_x, N_y) = (52, 28)$ as that used by CE95.

in many cases. To conclude this section, it is worth noting that the Milinazzo and Saffman[6] and CE95 homotopy (1) is only a particular choice of Poiseuille-Couette homotopy, and that more general transformations from one flow to the other, *e. g.*

$$U_B(y) = (1 - f(\eta))\left(1 - y^2\right) + g(\eta)\, y\ , \qquad (8)$$

where f and g satisfy the right end conditions, may lead to different results. This kind of behaviour has notably been reported in Taylor-Couette flow by Rucklidge and Champneys[16], who discovered that a connection between solutions for two different sets of boundary conditions was only possible in their problem for some special homotopy choices.

IV. SYMMETRY-PRESERVING HOMOTOPY

Motivated by the apparent similarities between the steady solutions of CE95 and those that MH05 obtained for small values of their perturbation parameter ε, we then considered the problem of finding steady two-dimensional nonlinear solutions in plane Couette flow ($\eta = 1$) using the homotopy proposed by MH05 and parameters similar to those used in their calculations. An ε symmetry-preserving identity perturbation was included in the equations for that purpose. As mentioned in Sec. II, this causes the least stable eigenvalue of the Couette flow to become unstable. For Re > 375 and $\alpha = 0.2$, the bifurcation is of the Hopf type, while for Re < 375 and the same α, the bifurcation is steady. In the latter case, for an identity perturbation with $\varepsilon > \varepsilon_0$, where $\varepsilon_0 \simeq 0.0456$, the eigenvalue crosses the imaginary axis for two values of Re (see MH05), leading to two different branches of steady nonlinear solutions (Fig. 4). MH05 showed that steady branches with $\varepsilon < \varepsilon_0$ cannot be computed directly because they reduce to a single point at $\varepsilon = \varepsilon_0$ and Re $\simeq 270$ (see their Fig. 11). Nevertheless, as noted in their paper, there is a connection between low Re branches with $\varepsilon > \varepsilon_0$ and high Re branches, including some with $\varepsilon < \varepsilon_0$. For instance, a solution at $\varepsilon = 0.04$ (dashed line) can be obtained by reducing ε, starting from a high energy solution at $\varepsilon = 0.05$ and Re $\simeq 360$. The associated $\varepsilon = 0.04$ steady branch can then be continued to large values of Re and transformed into high Re branches with different values of ε.

In Fig. 4, a travelling branch at $\varepsilon = 0.04547$ is also plotted. This branch connects to a steady high Re branch with the same ε at Re $= 500$. The presence of such a travelling branch, which was discovered by performing continuation with respect to ε in a range of Re where steady branches of solutions also exist, confirms the findings of MH05 that travelling solutions bifurcating at Re > 375 can turn into steady solutions by changing ε at fixed Re. Fig. 5 shows a typical continuation with respect to ε that clarifies this connection. Continuation in this figure is



FIG. 4: Bifurcations and nonlinear continuation of two-dimensional solutions of the ε-perturbed plane Couette flow of MH05 with $\alpha = 0.2$, $\varepsilon = 0.05$ (full line), $\varepsilon = 0.04$ (dashed line) and $\varepsilon = 0.04547$ (dotted and dash-dotted lines). $(N_x, N_y) = (32, 48)$ has been used (the results of some of these computations done at $(N_x, N_y) = (52, 48)$ are undistinguishable from those at $N_x = 32$). All solutions are steady except for the dash-dotted line branch at $\varepsilon = 0.04547$ (dash-dotted, red online), for which the phase speed reaches 0 when it meets the dotted steady branches. The vertical dash-triple dotted line corresponds to the path followed in the continuation presented in Fig. 5. Continuation was stopped at large energy (strong nonlinearity) due to insufficient numerical resolution.



FIG. 5: Nonlinear continuation of two-dimensional solutions of the ε-perturbed plane Couette flow of MH05 with respect to ε, for $\alpha = 0.2$, $\mathrm{Re} = 500$ and $(N_x, N_y) = (32, 48)$. The dashed line corresponds to the phase speed of the solutions and the full line to their kinetic energy. The starting point of the continuation corresponds to the Hopf bifurcation of the perturbed Couette flow, which occurs at $\varepsilon = 0.05358$ for $\mathrm{Re} = 500$. Continuation was stopped at large energy (strong nonlinearity) due to insufficient numerical resolution.



FIG. 6: Bifurcations and nonlinear continuation of steady two-dimensional $\alpha = 0.2$ solutions of the ε-perturbed plane Couette flow without the artifact mean flow perturbation for $\varepsilon = 0.05$ (full line), $\varepsilon = 0.05358$ (dashed line) and $\varepsilon = 0.04547$ (dash-dotted line) when the ε perturbation term is added to the $k_x \neq 0$ part of the equations only. $(N_x, N_y) = (32, 48)$. Continuation was stopped at large energy (strong nonlinearity) due to insufficient numerical resolution.

done along the line Re = 500 represented in Fig. 4. One finds that there are in fact two travelling branches of solutions with the same kinetic energy and opposite phase speed. These solutions merge at $\varepsilon = 0.04547$ and transform into a family of nonlinear steady solutions which corresponds to a cut at fixed Re of the surface of solutions in the $\mathrm{Re}-\varepsilon-$amplitude space. We conjecture that this behaviour is a reminiscence in the nonlinear regime of the collision between the first two real eigenvalues of the linear problem and their subsequent splitting into a complex conjugate pair (see MH05 for full details regarding the linear problem). Note finally that Fig. 5 is in qualitative agreement[18] with Fig. 6 of MH05.

The behaviour of steady solutions at large Re suggests that solutions with smaller and smaller values of ε can be found provided that Re is increased sufficiently. However, a careful examination of Fig. 4 reveals the presence of intriguing "double" bifurcations at large Re (see for instance the closed loop in dotted line emanating from the filled black circle at $\mathrm{Re} = 868.2$ and $\varepsilon = 0.04547$) that are not predicted by standard linear stability analysis for infinitesimal velocity perturbations with a single wavenumber $\alpha \neq 0$ (this also occurs in Fig. 12 of MH05). This behaviour can be explained as follows. Consider the equation for the perturbed mean flow

$$\mathrm{Re}\left(\partial_t \overline{u} + \partial_y \overline{uv}\right) = \partial_y^2 \overline{u} + \varepsilon \mathrm{Re} \overline{u}, \tag{9}$$

supplemented by the set of boundary conditions (3) on u. Eigenmodes of the linearized version of this equation



FIG. 7: Nonlinear continuation of steady $\alpha = 0.2$ two-dimensional solutions of the ε-perturbed plane Couette flow without the artifact mean flow perturbation as a function of ε, for Re = 1000 (full line), Re = 2500 (dotted line), Re = 9 000 (dashed line) and Re = 30 000 (dash-dotted line). $(N_x, N_y) = (32, 48)$.

take on the form

$$\overline{u} = A \sin(n\pi y), \tag{10}$$

and their growth rate γ is given by

$$\gamma = \varepsilon - (n\pi)^2/\mathrm{Re}\ . \tag{11}$$

These linear modes have no self-interactions, therefore they are also exact steady solutions of the fully nonlinear equations for any amplitude A when $\varepsilon\mathrm{Re} = (n\pi)^2$. For $n = 2$, we precisely obtain the bifurcation points observed in Fig. 4 at Re = 868.2 for $\varepsilon = 0.04547$ and at Re = 789.6 for $\varepsilon = 0.05$, which hints that these mean flow modes participate in the dynamics of high Re solutions described earlier. Note that the original idea of MH05 was to introduce an ε perturbation in order to cause the least stable eigenmode of Plane Couette flow with $k_x = \alpha$ to bifurcate, not to influence the whole dynamics of the flow including that of the $k_x = 0$ Fourier mode.

It is therefore natural to ask whether or not high Re, low ε branches survive when the artificial mean flow eigenmode is removed, and if solutions with $\varepsilon = 0$ can be found subsequently. Thanks to the way the problem has been set-up numerically in this study, it is straightforward to get rid of this model artefact by adding the ε perturbation in Eq. (4) but not in Eq. (5). Fig. 6 shows that branches of solutions at small Re computed with the corrected model look similar to those in Fig. 4, which were obtained with the original MH05 model. The main difference is for the lowest Re branch represented in these figures, which is supercritical in the MH05 model whereas it is subcritical when the mean flow perturbation is removed. A connection with high Re branches is also found in the corrected model, but in that case the branches do not bifurcate at some ε-dependent critical Re, even though they have a rather low kinetic energy at large Re. In order to discover nonlinear plane Couette flow solutions, we finally attempted to continue solutions obtained at a given Re with respect to ε. This continuation is presented in Fig. 7 for different values of Re. The amplitude of the solutions is seen to asymptote to infinity in the limit of vanishing ε for all selected values of Re. The natural conclusion is that no two-dimensional nonlinear solutions can be found for plane Couette flow using this homotopy technique.

V. CONCLUSIONS

The results presented in this study tend to show that one cannot identify steady nonlinear two-dimensional solutions in plane Couette flow using either the Poiseuille-Couette homotopy of CE95 or the ε-perturbation homotopy suggested by MH05. We have been unable to recover the solutions reported by CE95 and have pointed out that they may be spurious. It is therefore also likely that the results of Cherhabili and Ehrenstein[10] suffer from the same problem. Other more general homotopies like Eq. (8) may however be helpful to discover a connection between Poiseuille and Couette flow. The steady solutions obtained by MH05 at large Re and small ε values were shown to be influenced by an artifact of the perturbed equations, however using a revised version of their model did not help discovering new solutions for plane Couette flow because the corrected solutions found at small ε asymptote to infinite amplitude in the $\varepsilon = 0$ limit. This leaves the issue of the existence of two-dimensional nonlinear solutions in plane Couette flow unsettled. Another interesting avenue of investigation would be to consider the fate of the inviscid solutions discovered by Kida[17] in a uniform unbounded shear flow when a small amount of viscosity is added. The relevance of this type of solutions to wall-bounded flows is however quite unclear.

To conclude, it is worth pointing out that discovering nonlinear two-dimensional solutions in plane Couette flow is also relevant to the problem of the nonlinear stability of linearly stable swirling flows. Indeed, plane Couette flow with constant (spanwise) rotation can be used as a local model of differentially rotating flows. Since two-dimensional incompressible dynamics in plane Couette flow is not affected by the presence of a Coriolis force associated with a rotation vector perpendicular to the (x, y) plane, possible two-dimensional nonlinear solutions in non-rotating plane Couette flow would also be solutions in plane Couette flow with constant spanwise rotation for any rotation regime, including centrifugally stable anticyclonic regimes for which no purely hydrodynamic solutions are currently known at all[12].

Acknowledgments

The author acknowledges several helpful discussions with P. G. Mehta, G. I. Ogilvie and A. M. Rucklidge. This work has been supported by the Leverhulme Trust and the Newton Trust.

[1] K. H. Bech, N. Tillmark, P. H. Alfredsson, and H. I. Andersson. An investigation of turbulent plane couette flow at low reynolds numbers. *J. Fluid Mech.*, 286:291–325, 1995.

[2] M. Nagata. Three-dimensional finite-amplitude solutions in plane Couette flow: bifurcation from infinity. *J. Fluid Mech.*, 217:519–527, 1990.

[3] R. M. Clever and F. H. Busse. Three-dimensional convection in a horizontal fluid layer subjected to a constant shear. *J. Fluid Mech.*, 234:511–527, 1992.

[4] F. Waleffe. Exact coherent structures in channel flow. *J. Fluid Mech.*, 435:93–102, 2001.

[5] F. Waleffe. Homotopy of exact coherent structures in plane shear flows. *Phys. Fluids*, 15:1517–1534, 2003.

[6] F. A. Milinazzo and P. G. Saffman. Finite-amplitude steady waves in plane viscous shear flows. *J. Fluid Mech.*, 160:280, 1985.

[7] J. P. Zahn, J. Toomre, E. A. Spiegel, and D. O. Gough. Nonlinear cellular motions in Poiseuille flow. *J. Fluid Mech.*, 64:319–345, 1974.

[8] T. Herbert. Periodic secondary motions in a plane channel. In A. I. van de Vooren and P. J. Zandbergen, editors, *Proc. Int. Conf. Numer. Methods Fluid Dyn.*, pages 235–240. Berlin: Springer-Verlag, 1976.

[9] A. Cherhabili and U. Ehrenstein. Spatially localized two-dimensional finite-amplitude states in plane Couette flow. *Eur. J. Mech. B/Fluids*, 14:667, 1995.

[10] A. Cherhabili and U. Ehrenstein. Finite-amplitude equilibrium states in plane Couette flow. *J. Fluid Mech.*, 342: 159–177, 1997.

[11] P. G. Mehta and T. J. Healey. On steady solutions of symmetry-preserving perturbations of the two-dimensional Couette flow problem. *Phys. Fluids*, 17:4108, 2005.

[12] F. Rincon, G. I. Ogilvie, and C. Cossu. On self-sustaining processes in Rayleigh-stable rotating plane Couette flows and subcritical transition to turbulence in accretion disks. *Astron. & Astrophys.*, 463:817–832, 2007.

[13] H. Wedin and R. R. Kerswell. Exact coherent structures in pipe flow: travelling wave solutions. *J. Fluid Mech.*, 508:333–371, 2004.

[14] J. A. C. Weidemann and S. C. Reddy. A Matlab differentiation suite. *ACM Transactions on Mathematical Software*, 26(4):465–519, 2000.

[15] J. P. Boyd. *Chebyshev and Fourier Spectral Methods.* Dover, Mineola, New York, 2001. Second Edition.

[16] A. M. Rucklidge and A. R. Champneys. Boundary effects and the onset of Taylor vortices. *Physica D*, 191:282–296, 2004.

[17] S. Kida. Motion of an elliptic vortex in a uniform shear flow. *J. Phys. Soc. Japan*, 50:3517, 1981.

[18] In the course of this study, P. G. Mehta (private communication) pointed out to us that a lack of quantitative agreement is not surprising here due to the limited resolution capacities of the code used by MH05. Note that the code used in the present paper gives robust results even for very large systems of equations (including three-dimensional problems) and that it quantitatively reproduces the steady results of MH05 (compare Fig. 4 here and their Fig. 12).